

## Rapid Killing of *Mycobacterium terrae* by *N*-Chlorotaurine in the Presence of Ammonium is Caused by the Reaction Product Monochloramine

MARKUS NAGL AND WALDEMAR GOTTARDI

*Institute for Hygiene, Leopold-Franzens University of Innsbruck, Fritz-Pregl-Str. 3, A-6010 Innsbruck, Austria*

### Abstract

We have studied the activity of the weak endogenous oxidant *N*-chlorotaurine against *Mycobacterium terrae*.

The study revealed slow killing of more than 2 h duration by 1% (55 mM) *N*-chlorotaurine. In the presence of ammonium chloride, however, killing times decreased to a few minutes, even by 0.1% *N*-chlorotaurine.

This phenomenon is explained by formation of the lipophilic and therefore more bactericidal monochloramine as a result of transhalogenation of ammonia by *N*-chlorotaurine.

*N*-Chlorotaurine (Cl-HN-CH<sub>2</sub>-CH<sub>2</sub>-SO<sub>3</sub>H), the *N*-chloro derivative of the amino acid taurine, is a weak oxidant produced at high concentrations in man by granulocytes and monocytes (Lampert & Weiss 1983; Grisham et al 1984). *N*-Chlorotaurine has significant microbicidal efficacy against bacteria, yeasts (Nagl & Gottardi 1992, 1996) viruses (Nagl et al 1998a) and worms (Yazdanbakhsh et al 1987).

The bactericidal and fungicidal activity of *N*-chlorotaurine has been shown to be enhanced by the presence of low molecular-weight amino compounds and also in inflammation samples because of transformation to more bactericidal *N*-chloro derivatives (Nagl & Gottardi 1996). Rapid inactivation of omniresistant *Pseudomonas aeruginosa* by *N*-chlorotaurine in urine has been considered to be connected with the formation of monochloramine (NH<sub>2</sub>Cl) by halogenation of ammonium (Nagl et al 1998c). The bactericidal activity of NH<sub>2</sub>Cl has already been proved (LeChevallier et al 1988; Jacangelo et al 1991). Its mycobactericidal effect, however, has not yet been investigated.

This study was performed to examine whether the in-situ production of monochloramine by reaction of ammonium and *N*-chlorotaurine could induce sufficient activity against mycobacteria which are usually highly resistant to disinfectants.

Correspondence: M. Nagl, Institute for Hygiene, Leopold-Franzens University of Innsbruck, Fritz-Pregl-Str. 3, A-6010 Innsbruck, Austria.

To verify that the bactericidal effect is caused solely by NH<sub>2</sub>Cl and not by other *N*-chloro compounds of low molecular weight (amino acids and oligopeptides from the agar), the experiments were conducted with thoroughly washed bacteria.

### Materials and Methods

#### Reagents

Pure *N*-chlorotaurine as the crystalline sodium salt (MW 181.52 g mol<sup>-1</sup>) (Nagl & Gottardi 1996) was dissolved in 0.1 M phosphate buffer, pH 6.8–7.2. The identity of the *N*-chlorotaurine was proved by IR spectrometry, purity by iodometric titration (calculated 19.53% Cl<sup>+</sup>, found 19.3% Cl<sup>+</sup>, purity 99%). Reagent grade ammonium chloride, sodium thiosulphate and buffers (sodium dihydrogen phosphate and disodium hydrogen phosphate) were purchased from Merck (Darmstadt, Germany).

*Bacteria, media and assay of antibacterial action*  
*Mycobacterium terrae* ATCC 15755 (van Klingeren Pullen 1987; Sattar et al 1995) was obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) and grown on Löwenstein-Jensen slant agar (Becton Dickinson, USA) at 37°C for 21 days until a layer of non-countable colonies had formed. After addition of NaCl (0.9%, w/v; 10 mL) to the agar tube, bacteria were removed from the agar by sonication

for 2 min (Bandelin Sonorex RK 102 water bath, 35 kHz, 120/240 W) without loss of viability. Bacteria were centrifuged twice at 1800 *g* for 10 min and washed twice with NaCl (0.9%) to remove from the suspension organic compounds which could have reacted with *N*-chlorotaurine and falsified the results. The suspensions were vortex-mixed vigorously ( $3 \times 5$  s) before each centrifugation and sonicated again for 2 min to furnish a homogenous suspension of  $2-9 \times 10^7$  colony-forming units  $\text{mL}^{-1}$ . This suspension (0.1 mL) was added to solutions (0.9 mL) of *N*-chlorotaurine or of *N*-chlorotaurine +  $\text{NH}_4\text{Cl}$ . Controls without *N*-chlorotaurine were treated the same way.

Subsequent to incubation for 1, 5, 10 and 20 min at room temperature, samples (0.1 mL) were mixed with sodium thiosulphate solution (3–9%, w/v; 0.1 mL) to inactivate 0.1%, 1% and 3% (w/v) *N*-chlorotaurine, respectively. This solution (50  $\mu\text{L}$ ) and similar volumes of serial tenfold dilutions in 0.9% NaCl were added to Löwenstein-Jensen slant agar which was carefully tilted five times to distribute the bacteria optimally on the agar surface and then stored at 37°C for 6 weeks. Colony-forming units were counted after 3 weeks of growth, later evaluations did not reveal any change in the results.

## Results

### Action of *N*-chlorotaurine without additives

In buffer solution, no killing by 1% *N*-chlorotaurine was observed within 120 min incubation time. Incubation times of 7 days and 14 days yielded a reduction in colony-forming units of  $4.0 \pm 1.2$  and  $5.9 \pm 0.5 \log_{10}$ , respectively, for samples containing 1% *N*-chlorotaurine and of  $1.0 \pm 0.5$  and  $1.5 \pm 0.4 \log_{10}$ , respectively, for controls (mean values  $\pm$  standard deviation,  $n=5$ ,  $P < 0.01$  by use of Student's paired *t*-test).

### Action of *N*-chlorotaurine in the presence of ammonium chloride

In contrast with *N*-chlorotaurine without additives, addition of ammonium chloride resulted in significant bactericidal activity after incubation times of only a few minutes. Time-kill curves are shown in Figure 1. Killing proved to be more dependent on the concentration of ammonium chloride than on that of *N*-chlorotaurine.

As expected, no reduction in the number of colony-forming units could be observed in presence of 0.1–10% ammonium chloride without *N*-chlorotaurine. Inactivation of 10% ammonium chloride and 3% *N*-chlorotaurine by addition of 9% sodium thiosulphate before addition of the bacteria also removed any bactericidal effect.




Figure 1. Killing of *M. terrae* by 0.1% (upper), 1.0% (middle) and 3.0% (lower) *N*-chlorotaurine in the presence of 0.1% ( $\square$ ), 1% ( $\triangle$ ) or 10% ( $\times$ )  $\text{NH}_4\text{Cl}$  at room temperature and pH 6.8–7.2 (7.8 for 3% *N*-chlorotaurine + 0.1%  $\text{NH}_4\text{Cl}$ ). Results from controls ( $\blacklozenge$ ) in phosphate buffer without any additives were no different from those containing 10%  $\text{NH}_4\text{Cl}$  only. Values are means  $\pm$  standard deviations of colony-forming units ( $\log_{10}$ )  $\text{mL}^{-1}$ ,  $n=3-6$ ;  $P < 0.01$  for all cases with  $\geq 0.5 \log_{10}$  difference between samples and controls (Student's paired *t*-test).

The very large increase in killing power in the presence of ammonium chloride can, therefore, be attributed solely to the formation of  $\text{NH}_2\text{Cl}$ , which proved to be a highly mycobactericidal agent.

## Discussion


In this study the mycobactericidal action of *N*-chlorotaurine has been investigated for the first time. Although, as expected, the action proved to be weaker than that against other Gram-positive and Gram-negative bacteria (Nagl & Gottardi 1992,

Table 1. Equilibrium concentrations\* of NH<sub>2</sub>Cl (mM) in aqueous solutions from different initial concentrations of ammonium chloride and *N*-chlorotaurine at pH 7.0.

Concentration of <i>N</i> -chlorotaurine (% (mM))	Concentration of ammonium chloride (% and (mM))		
	0.1 (18.7)	1.0 (187.0)	10.0 (1869.5)
3.0 (165.3)	4.7	14.7	34.6
1.0 (55.1)	3.1	8.7	18.5
0.1 (5.5)	1.1	2.4	3.9

\* Calculated with  $K_{\text{NH}_2\text{Cl}} = 0.02$ .

1996), it was improved to a surprising extent by ammonium chloride, which produces monochloramine (NH<sub>2</sub>Cl) by transhalogenation (Thomas 1979; Grisham et al 1984; Nagl & Gottardi 1996):



Increasing the molar ratio NH<sub>4</sub>Cl/*N*-chlorotaurine significantly increased the speed of mycobactericidal action, indicating increased formation of NH<sub>2</sub>Cl, in accordance with the law of mass action. With the known equilibrium constant of equation 1 ( $K_{\text{NH}_2\text{Cl}} = 0.02$ ; Nagl & Gottardi 1996) the concentration of NH<sub>2</sub>Cl in these experiments could be estimated (Table 1). It is apparent that there is a good correlation between calculated NH<sub>2</sub>Cl concentrations and bactericidal activity denoted as log<sub>10</sub> reduction in colony-forming units. From nine results with 1-min incubation time linear regression ( $R = 0.91$ ) gave an average of  $0.00497 \pm 0.00080 \text{ mol L}^{-1}$  NH<sub>2</sub>Cl necessary to effect a one log<sub>10</sub> kill.

Because mycobacteria are, because of their special cell wall with high lipid content (Russell et al 1986), more resistant to chemical attack than other vegetative bacteria, NH<sub>2</sub>Cl as a lipophilic compound (Grisham et al 1984) will be able to penetrate mycobacteria more easily than the hydrophilic *N*-chlorotaurine. Because the substantial difference between the action of *N*-chlorotaurine and NH<sub>2</sub>Cl is unrelated to reactivity, but more to size and polarity (Thomas 1979; Nagl & Gottardi 1996), our findings accentuate the importance of penetration for rapid mycobactericidal action. This consideration is in keeping with the mechanism of inactivation of *Escherichia coli* by NH<sub>2</sub>Cl, which is based on inhibition of respiration, substrate dehydrogenation and bacterial transport rather than on damage to the cell envelope (Jacangelo et al 1991).

This study reveals the importance of NH<sub>2</sub>Cl as a highly effective bactericidal substance. However, because of its low stability it cannot be stored and has to be produced in-situ from ammonia by a chlorinating agent. For this reason *N*-chlorotaurine, which is well tolerated by living tissue (Nagl et al 1998b, c), must be preferred to other, more strongly oxidizing and therefore more irritating compounds such as chloramine T or hypochlorite.

Further studies will be necessary to evaluate the role of endogenous ammonium which obviously acts as an amplifier of the bactericidal properties of chlorine-based oxidants.

#### Acknowledgements

We wish to thank Professor Dr Manfred P. Dierich, head of the department, for continuous support. Since July 1997 this work has been supported by grant no. P12298-MED of the FWF (Austrian Science Fund).

#### References

- Grisham, M. B., Jefferson, M. M., Melton, D. F., Thomas, E. L. (1984) Chlorination of endogenous amines by isolated neutrophils. *J. Biol. Chem.* 259: 10404–10413
- Jacangelo, J. G., Olivieri, V. P., Kawata, K. (1991) Investigating the mechanism of inactivation of *Escherichia coli* B by monochloramine. *J. AWWA* 83: 80–87
- Lampert, M. B., Weiss, S. J. (1983) The chlorinating potential of the human monocyte. *Blood* 62: 645–651
- LeChevallier, M. W., Cawthon, C. D., Lee, R. G. (1988) Inactivation of biofilm bacteria. *Appl. Environ. Microbiol.* 54: 2492–2499
- Nagl, M., Gottardi, W. (1992) In vitro experiments on the bactericidal action of *N*-chlorotaurine. *Hyg. Med.* 17: 431–439
- Nagl, M., Gottardi, W. (1996) Enhancement of the bactericidal efficacy of *N*-chlorotaurine by inflammation samples and selected N-H compounds. *Hyg. Med.* 21: 597–605
- Nagl, M., Larcher, C., Gottardi, W. (1998a) Activity of *N*-chlorotaurine against herpes simplex- and adenoviruses. *Antiviral Res.* 38: 25–30

- Nagl, M., Miller, B., Daxecker, F., Ulmer, H., Gottardi, W. (1998b) Tolerance of *N*-chlorotaurine, an endogenous antimicrobial agent, in the rabbit and human eye—a phase I clinical study. *J. Ocular Pharmacol. Ther.* 14: 283–290
- Nagl, M., Pfausler, B., Schmutzhard, E., Fille, M., Gottardi, W. (1998c) Tolerance and bactericidal action of *N*-chlorotaurine in a urinary tract infection by an omniresistant *Pseudomonas aeruginosa*. *Zbl. Bakt.* In press
- Russell, A. D., Hammond, S., Morgan, J. R. (1986) Bacterial resistance to antiseptics and disinfectants. *J. Hosp. Infect.* 7: 213–235
- Sattar, S. A., Best, M., Sprinthorpe, V. S., Sanani, G. (1995) Mycobactericidal testing of disinfectants: an update. *J. Hosp. Infect.* 30 (Suppl): 372–382
- Thomas, E. L. (1979) Myeloperoxidase, hydrogen peroxide, chloride antimicrobial system: effect of exogenous amines on the antibacterial action against *Escherichia coli*. *Infect. Immun.* 25: 110–116
- van Klingeren, B., Pullen, W. (1987) Comparative testing of disinfectants against *Mycobacterium tuberculosis* and *Mycobacterium terrae* in a quantitative suspension test. *J. Hosp. Infect.* 10: 292–298
- Yazdanbakhsh, M., Eckmann, C. M., Roos, D. (1987) Killing of schistosomula by taurine chloramine and taurine bromamine. *Am. J. Trop. Med. Hyg.* 37: 106–110